

SCH 23390: The First Selective Dopamine D₁-Like Receptor Antagonist

James A. Bourne

*Department of Physiology, Monash University,
Clayton, Victoria, Australia*

Key Words: Dopamine—Dopamine D₁ receptors—Epileptiform activity—Neurological disorders—SCH 23390.

ABSTRACT

SCH 23390, the halobenzazepine (*R*)-(+)-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine, is a highly potent and selective dopamine D₁-like receptor antagonist with a K_i of 0.2 and 0.3 nM for the D₁ and D₅ dopamine receptor subtypes, respectively. *In vitro*, it also binds with high affinity to the 5-HT₂ and 5-HT_{1C} serotonin receptor subtypes. However, the doses required to induce a similar response *in vivo* are greater than 10-fold higher than those required to induce a D₁-mediated response.

Previous *in vivo* pharmacological studies with SCH 23390 have shown it to abolish generalized seizures evoked by the chemoconvulsants: pilocarpine and soman. These studies provide evidence of the potential importance of D₁-like dopaminergic receptor mechanisms in facilitating the initiation and spread of seizures. The inference from a majority of studies is that the activation of dopamine D₁ receptors facilitates seizure activity, whereas activation of D₂ receptors may inhibit the development of seizures. SCH 23390 has also been used in studies of other neurological disorders in which the dopamine system has been implicated, such as psychosis and Parkinson's disease. Apart from the study of neurological disorders, SCH 23390 has been extensively used as a tool in the topographical determination of brain D₁ receptors in rodents, nonhuman primates, and humans.

In summary, SCH 23390 has been a major tool in gaining a better understanding of the role of the dopamine system, more specifically the D₁ receptor, in neurological function and dysfunction.

Address correspondence and reprint requests to: Dr. James A. Bourne, Department of Physiology, Monash University, Clayton, Victoria, 3800, Australia. E-mail: James.bourne@med.monash.edu.au.

INTRODUCTION

Dopamine receptors have been associated with many neurological disorders, including most notably Parkinson's disease and schizophrenia. Pharmacological studies carried out before the advent of receptor cloning techniques supported the existence of just two dopamine receptors, termed D₁ and D₂, based on their positive and negative coupling to adenylate cyclase, mediated by a G-protein and effector (45). For example, D₁ receptor activation enhances cAMP formation, whereas activation of D₂ receptors inhibits cAMP formation. However, it was not until the 1980s that cloning studies revealed that there are at least five receptors, which have been termed D₁–D₅. The cloning of the D₂ splice variants (D_{2L} and D_{2S}) was followed by the cloning of products of different genes, such as the D₃ (60), D₄ (68), and D₅ receptors (64). These receptor subtypes were then assigned to either the D₁-like subfamily (which includes the D₁ and D₅ receptors) or D₂-like subfamily (which includes the D₂, D₃, and D₄ receptors), depending on their coupling to adenylate cyclase. Analysis of the amino acid sequences of the dopamine receptor subtypes reveals that there is a high degree of homology between the members of the subfamilies (43). The genomic organization of the dopamine receptors also supports the notion that they derive from the divergence of two gene subfamilies (49).

The D₁-like receptor proteins, similar to the D₂-like receptors, contain seven hydrophobic alpha-helical transmembrane spanning segments linked to more hydrophilic segments with an extracellular amino terminus. This is the classical conformation for G-protein-coupled receptors (28).

D₁ receptors are found in high levels in the dopamine rich areas of the brain, such as the caudate nucleus, putamen, substantia nigra, nucleus accumbens, hypothalamus, thalamus, frontal cortex, and olfactory tubercle, whereas the D₅ receptor is present, although at a low concentrations, in the hippocampus, thalamus, lateral mammillary nucleus, and cerebral cortex (35). Both of these receptor subtypes are capable of stimulating adenylyl cyclase (65). Gene transcripts of D₃, D₄, and D₅ receptors are much less abundant and display more discrete expression in the brain than those of the D₁ (59). In contrast, those of D₁ and D₂ are virtually absent from the dorsal striatum, which are only expressed in certain limbic regions (e.g., hippocampus). A consistent difference between the D₁ and D₅ receptors is that dopamine is approximately 10 times more potent at the D₅ receptor (Table 1). The function of the D₅ receptor is not understood but the D₁ receptor seems to mediate important central nervous system (CNS) actions of dopamine to control movement (17) and cognitive function (55).

SCH 23390 is a potent, enantioselective dopamine D₁-like receptor antagonist (11,40,41,53), although *in vivo* studies suggest that SCH 23390 may also have a moderately high affinity for the 5-HT₂ and 5-HT_{1C} receptors (12). The affinity of ligands at dopamine receptors is summarized in Table 1. SCH 23390 is a very short-acting compound with an elimination half-life of around 25 min following administration of 0.3 mg/kg i.p. in the rat (46). The few pharmacokinetic data available suggest that the compound undergoes extensive first-pass metabolism when administered orally (41).

Over the last two decades, our understanding of the function of the dopamine receptor and its neuronal location has been much aided by this drug. For example, it has been a widely used tool in behavioral and radioligand binding studies. In addition, more recent studies have proposed its effectiveness as an anticonvulsant in certain experimental paradigms.




Fig. 1. Chemical structure of SCH 23390, (*R*)-(+)-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine.

CHEMISTRY

SCH 23390, the 3-methyl, 7-chloro analogue of the D₁ agonist SKF 38393, is a potent enantioselective dopamine D₁-like receptor antagonist. The active enantiomer of the benzazepine ligand (C₁₇H₁₈ClNO) is (*R*)-(+)-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine (Fig. 1), compared with the inactive (*S*) enantiomer (SCH 23388). The commercially available hydrochloride salt has a molecular weight of 324.25 and is a white solid soluble in water (8 mg/mL), DMSO (3 mg/mL), or ethanol (2 mg/mL). Dopamine D₁-like receptor antagonists are commonly 1-phenyl-3-methyl-7-halogenated- benzazepines. Both the 3-methyl and 7-chloro substituents impart antagonist activity, however, the halogen is also essential (42). Also, an important part of

TABLE 1. Dissociation constants (K_i) for some of the selective/unselective dopamine antagonists/agonists at dopamine receptors

Ligand	K _i values (nm)				
	D ₁ -like		D ₂ -like		
	D ₁	D ₅	D ₂	D ₃	D ₄
Antagonists					
Chlorpromazine	~90	~130	3	4	35
Clozapine	~170	~330	~230	~170	21
Haloperidol	~80	~100	1.2	~7	2.3
SCH 23390	~0.2	0.3	~1100	~800	~3000
S-Sulpiride	~45,000	~77,000	~15	~13	1000
R-Sulpiride	~19,000	29,000	~900	~400	970
Agonists					
(-)-Apomorphine	~0.7		~0.7	~32	~4
Dopamine	0.9	<0.9	~7	~4	~30
7-OH-DPAT	~5000		10	~1	650
SKF 38393	1	~0.5	~150	~5000	~1000

the interaction between the phenyl ring in the benzazepines and the receptor is due to electrostatic forces with the phenyl ring interacting with the same receptor site as the oxygen atom of the 8-hydroxy group (54).

NEUROCHEMISTRY

Studies of changes in brain neurochemistry following the administration of SCH 23390 *in vivo* have focused primarily on the dopamine-rich nuclei, including the caudate, putamen, and accumbens, and have primarily utilized microdialysis (73,74) to monitor changes. In one *in vivo* study, local perfusion (90 min) of SCH 23390 (1 and 10 μ M) into the striatum of halothane-anesthetized rats increased extracellular dopamine over basal levels dose dependently, but had no effect on gamma-aminobutyric acid (GABA) and glutamate levels (50). The increase in striatal extracellular dopamine in an anesthetized preparation was not observed following parenteral or intranigral infusion (57,77). However, a dose-dependent increase in striatal extracellular dopamine and its metabolites dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA) was observed by Imperato et al. (1987) in the unanesthetized rat, following doses greater than 0.012 mg/kg (subcutaneous [s.c.]). Tomiyama et al. (66) suggest that the SCH 23390-evoked increase in dopamine and its metabolites is mediated by the reverse operation of the dopamine transporter, and not the D₁ receptor, because of the insensitivity of SCH 23390 to TTX, a sodium-channel blocker. The lack of increase in striatal extracellular GABA from basal levels is probably due to the inhibition of the dopamine D₁ release-enhancing heteroreceptors located on the terminals of the GABAergic interneurons (56). A decrease in acetylcholine and choline was also observed in the rat striatum after infusion of SCH 23390, probably due to a decrease in the release and turnover of acetylcholine, that originate from the striatal cholinergic interneurons (29). When studying the effects of drugs on neurotransmitters in an awake animal, the effect of environmental stress in these and other studies has been shown to alter the extracellular concentration of neurotransmitters in many brain locations. Studies have conclusively linked environmental stress with increases in the levels of dopamine and other neurotransmitters in the brain (27,31).

More recent *in vivo* studies have been directed at the prefrontal cortex (PFC), where the A10 dopaminergic neurons, which originate in the ventral tegmental area, project (6). Perfusion with the D₁ agonist SKF 38393 (2–200 μ M), via a dialysis probe into the medial PFC dose-relatedly reduced extracellular concentrations of both glutamate and GABA. The neurochemical changes evoked by 200 μ M SKF 38393 were, however, prevented by co-perfusion with SCH 23390 (40 μ M). These results suggest that the dopaminergic hyperactivity may lead to the hypofunction of glutamatergic and GABAergic systems in the PFC via D₁ dopamine receptor stimulation (1).

In one *in vitro* study, striatal slices from the rat were preincubated with [³H]GABA and superfused in the presence of nipecotic and aminooxyacetic acids, which are inhibitors of high-affinity GABA transport and GABA aminotransferase, respectively. GABA efflux was estimated by monitoring tritium efflux, 98% of which was in the form of [³H]GABA. The overflow of GABA evoked by electrical field stimulation (8 Hz) was increased two-fold by SKF 38393 (10 μ M) but was completely blocked by SCH 23390 (10 μ M). However, SCH 23390 had no effect on GABA overflow when given alone. Thus, activation of

dopamine D₁ receptors appears to exert an excitatory influence on GABA release; however, this effect was not elicited by endogenous dopamine under the conditions of this experiment (36). Another *in vitro* study examined the effects of SCH 23390 on the spontaneous and calcium-dependent, potassium-induced release of [³H]GABA accumulated by slices of rat substantia nigra. SKF 38393 and dopamine were without effect on [³H]GABA efflux when administered by themselves (1–40 μM) or in the presence of the phosphodiesterase inhibitor isobutylmethylxanthine (IBMX) (0.5 μM), but they potentiated evoked release in the presence of forskolin (0.5 μM), an adenylate cyclase activator (62).

PHARMACOLOGY

Effects on Motor Behavior

The pharmacologic profile of SCH 23390 resembles that of many other neuroleptics, and early studies showed it had an antistereotypic effect in mice, rats, and dogs, as well as a cataleptogenic effect and an inhibitory effect on amphetamine-induced circling in rats (21,22). In rodents, SCH 23390 inhibited motor stimulation induced by apomorphine or by selective D₂ agonists, such as RU 24213. This inhibition is evident across a range of behaviors (26,61), and it has been suggested that SCH 23390 blocks all motor activity (61). This result suggests that there is a definite synergy/cooperation between the D₁ and D₂ receptor subtypes in motor behavior. However, in drug discrimination paradigms SCH 23390 blocked the discriminative stimulus properties of amphetamine but not apomorphine (71), suggesting that the two receptor subtypes can act in a functionally independent manner to regulate response. Studies with SCH 23390 in nociceptive and thermic paradigms have shown that it can reverse the increase in tail flick latency evoked by the D₁ agonist SKF 38393, following long-term treatment with haloperidol (9).

More recently studies with SCH 23390 have focused on understanding the role of the basal ganglia in motor behavior to gain a better insight into the role of specific dopamine receptors in diseases such as idiopathic parkinsonism. The role of dopamine in the subthalamic nucleus to control motor behavior was investigated in rats using bilateral microinfusions of SCH 23390, which induced catalepsy. These findings suggest that dopamine D₁ receptors within the subthalamic nucleus (STN) play an important role in the regulation of motor functions (38). In the globus pallidus (GP) of rats bilateral microinfusions of SCH 23390 were used to investigate the role of dopamine receptors in motor control. Results show that SCH 23390 injected into the GP induced akinesia, as determined by means of the catalepsy test. These findings indicate that pallidal dopamine D₁ receptors are critically involved in the control of motor behavior. These findings further imply that defective dopaminergic transmission in the GP and STN might contribute to akinesia due to lesion- or drug-induced dopamine hypofunction in experimental animals and in neurodegenerative diseases (37).

Effects on Memory

Recent studies investigating the role of dopamine in working memory suggest that supranormal stimulation of D₁ receptors may contribute to the detrimental actions of do-

pamine on spatial working memory in the prefrontal cortex (8,51). One study directly tested this hypothesis by examining the effects of bilateral intra-PFC administration of the D₁ receptor agonist SKF 81297 (0.1 µg) into rats performing the spatial working memory task (T-maze). SKF 81297 produced a dose-related impairment in a working memory task. The impairment was reversed by pretreatment with SCH 23390 (0.01 mg/mL, i.p.). SCH 23390 by itself had no effect on performance cortex (8,51), demonstrating that supranormal D₁ receptor stimulation in the PFC is sufficient to impair PFC working function. These data are consistent with recent electrophysiological studies of D₁ receptor mechanisms affecting the PFC (75,76). However, recently Seamans et al. (58) have shown that local infusion of SCH 23390 (0.05–5 µg) into the prefrontal cortex blocked delayed-response task in rats on a radial-arm maze. Attentional performance was also tested using a five-choice serial reaction time task following intracortical (prefrontal) injection of SCH 23390 (0.3 µg), and selectively impaired the accuracy in high baseline correct responses (34). Conversely, intracortical administration of the D₁ receptor agonist SKF 38393 (0.6 µg) improved the accuracy, as well as the speed in making accurate responses, in the performance of the same task in rats that have low baseline correct responses. The latter improvement was antagonized by intracortical administration of SCH 23390 (1 µg) (34).

***In Vivo* Anticonvulsant Studies**

The predominantly proconvulsant action of the nonselective antipsychotic drugs in humans first indicated that dopaminergic pathways might play a crucial physiological role in suppressing the genesis and dissemination of seizure activity in the brain (63). Since then, the generation of specific dopamine receptor antagonists and agonists has enabled much work into the involvement of the subtypes of dopamine receptor in epilepsy. Many models of epileptiform activity have been used to investigate the role of specific receptor subtypes in seizure activity. The commonly used species include rats, mice, gerbils, and rabbits, with little attention paid to the guinea pig. The methods of inducing seizures have included the use of genetically prone animals, kindling, electroshock, and the use of chemoconvulsants such as strychnine, soman, pentylenetetrazol (PTZ), and pilocarpine, among others. More recently, the advent of new technologies has made it possible to concomitantly study the behavioral, neurochemical, and electrophysiological effects of seizure activity *in vivo* by means of an assembly based upon the conventional microdialysis probe (13).

Behavioral Studies

The pilocarpine (muscarinic acetylcholine receptor agonist) model in rats and mice has been extensively used and is a reliable model for studying the processes of secondary generalized seizures initiated in the hippocampus. Previous studies have reported that the D₁ antagonist SCH 23390 (pretreatment with 0.1–0.3 mg/kg, i.p., in rats) dose-dependently abolished seizures evoked by a low dose of pilocarpine (200 mg/kg, i.p.) (10). However, motor seizures evoked by a higher dose of pilocarpine (600 mg/kg) were more difficult to suppress, and Turski et al. (67) reported that SCH 23390 was completely ineffective in this situation. However, Al-Tajir and Starr (2,4) showed that the intrastriatal injection of SCH 23390 (1 µg in each striatum) was capable of delaying the onset of forelimb myoclonus (repeated muscle contractions) and reduced the severity and lethality of motor (con-

vulsive) seizures evoked by pilocarpine (600 mg/kg, i.p.). This study also highlighted the previously held assumption that the dopamine receptor agonists/antagonists are more effective following their direct application to the brain. Burke et al. (16) showed that pretreatment of mice with SCH 23390 (0.8 mg/kg, i.p.), 60 min prior to pilocarpine 400 mg/kg, prevented fatalities and only 14.3% of animals exhibited convulsions, a proportion that was significantly lower than seen with pilocarpine treatment alone. More recently, pretreatment with SCH 23390 (0.5 mg/kg, i.p.) has been shown to prevent the onset of the predefined clinical signs associated with soman-evoked seizure activity (30.5 µg/kg, s.c.) in the guinea pig (Table 2), suggesting the involvement of the D₁ receptor subtype in the manifestation of these behaviors, probably through activation of the striatonigral GABAergic pathway (14). SCH 23390 has also afforded protection against pilocarpine-induced convulsions following the intrastriatal and intraaccumbal administration of the antagonist (2), but not following intrahippocampal administration (where it only delayed the onset of "limbic motor seizures") (5). These findings provide evidence of the involvement of the basal ganglia in the manifestation of motor convulsions evoked by cholinergic mechanisms. Stimulation of the D₂ receptor, following intranigral administration of the D₂ agonist LY 171555, did not protect against pilocarpine-induced convulsions (3). SCH 23390 at 0.3 and 0.5 mg/kg was, however, ineffective at preventing PTZ-evoked seizures in mice.

Neurochemical Studies

In the soman model (30.5 µg/kg, s.c.) of seizure activity in the guinea pig, pretreatment of SCH 23390 (0.5 mg/kg, i.p.) resulted in changes to the striatal neurochemistry. Inhibition of the dopamine D₁ receptor evoked a relative decrease in striatal extracellular dopamine and its metabolites (HVA and DOPAC) compared with that seen with soman alone, in which the animal had electrographic and behavioral seizures (Fig. 2). This suggests that the level of striatal extracellular dopamine is strongly influenced by the seizure activity evoked by the chemoconvulsant soman. Also in the same study, a relative decrease was observed in the concentration of striatal extracellular acetylcholine (ACh), suggesting that the D₁ heteroreceptors located on the striatal intrinsic cholinergic neurons are capable of modulating the release of ACh in an excitatory manner (14). In guinea pigs

TABLE 2. Time to onset (min; mean ± S.E.M., n = 5) of pre-defined clinical signs, following the administration of soman (30.5 µg/kg). None of these clinical signs were observed in animals pre-treated with SCH 23390 (0.5 mg/kg)

Clinical sign	Time to onset (min)
Chewing	12 ± 1
Haunched posture	10 ± 3
Splayed hind limbs	27 ± 4
Circling	28 ± 4
Tremor	32 ± 4
Writhing, ataxia	56 ± 11
Myoclonic convulsions	53 ± 11
Vocalization	79 ± 9




Fig. 2. Evoked changes “total effect” in the striatal extracellular dialysate content of dopamine, DOPAC and HVA following administration of soman (30.5 µg/kg; solid bars) and SCH 23390 (0.5 µg/kg) + soman (open bars), in the guinea pig. Results are expressed as a percent of the respective baseline values (mean ± S.E.M., $n = 5$). ** $P < 0.01$; *** $P < 0.001$, soman vs. SCH 23390 + soman; two-tailed Mann–Whitney U -test.

treated with the excitotoxin, kainic acid (12 mg/kg), and pretreated with SCH 23390 (0.5 mg/kg, i.p.) there was incomplete attenuation of seizure activity; this was accompanied by a significant relative decrease in dopamine, but not its metabolites (Fig. 3). A decrease was observed also in the striatal extracellular aspartate and glutamate levels. However, an increase in GABA levels was observed (Fig. 4) (15). The increase in dopamine, compared with that observed with kainic acid alone, could be a result of the activation of kainic acid release-enhancing heteroreceptors, located on the terminals of the nigrostriatal dopaminergic neurons (47,48). The lack of change in the metabolites DOPAC and HVA suggests no increase in the metabolism of dopamine. The decrease in the striatal extracellular concentration of excitatory amino acid neurotransmitters aspartate and glutamate, compared with that seen with kainic acid alone, suggests a decrease in the activation of the corticostriatal glutamatergic fibers, presumably due to a decrease in the activation of the cortex (seen in the EEG). The increase in GABA is more difficult to explain, since the inhibition of the dopamine D_1 heteroreceptors located on the striatal GABAergic interneurons would usually result in a decrease in the release of GABA (8,51). Therefore, other receptors or pathways may be implicated.

Electrophysiological Studies

The capability of the D_1 antagonist SCH 23390 to prevent cortical electrographic (ECoG) paroxysmal activity has been observed in the pilocarpine model (200 mg/kg, i.p.) of epileptiform activity in rats (10). This specific activity of the dopamine D_1 receptor was also supported by results in the same study, which showed that systemic administration of the D_1 agonist SKF 38393 exacerbated the seizure activity relative to that seen with higher




Fig. 3. Evoked changes “total effect” in the striatal extracellular dialysate content of dopamine, DOPAC and HVA following administration of kainic acid (12 mg/kg; solid bars) and SCH 23390 (0.5 mg/kg) + kainic acid (open bars), in the guinea pig. Results are expressed as a percent of the respective baseline values (mean \pm S.E.M., $n = 5$). *** $P < 0.001$, kainic acid vs. SCH 23390 + kainic acid; two-tailed Mann–Whitney U -test.

doses of the chemoconvulsant. In the soman model (30.5 $\mu\text{g/kg}$, s.c.) of seizure activity in the guinea pig, pretreatment with SCH 23390 (0.5 mg/kg, i.p.) was capable of preventing the onset of electrographic seizure activity in both the striatum (ESTG) and the cortex. This resulted in a significant decrease in the EEG power compared with that seen with soman alone (Fig. 5), which was at a similar level to that seen prior to the injection of the chemoconvulsant. Conversely, following pretreatment with the D_2 receptor antagonist sulpiride (30 mg/kg), there was an earlier onset of an EEG-defined *status epilepticus* and an increase in the power of the EEG but not in behavioral paroxysmal activity (14). With kainic acid (12 mg/kg, i.p.) and pretreatment with SCH 23390 (0.5 mg/kg, i.p.) the dopamine antagonist did not totally ameliorate the paroxysmal activity but reduced the severity of the seizure activity in terms of the power of the EEG (Fig. 6), and more interictal periods were observed (15). The D_2 receptor agonist LY 171555 (0.5 mg/kg) similarly depressed pilocarpine-induced seizures when administered subcutaneously (4). These studies indicate that the two dopamine receptor subtypes, D_1 and D_2 , exert opposing roles in the generation of electrographic seizure activity. While it may be an overgeneralization to characterize D_1 and D_2 receptors as mediating pro- and anticonvulsive roles, empirical evidence certainly characterizes both a detrimental and beneficial function in man.

One caveat that has come from these studies is the importance of distinguishing electrographic and behavioral seizure activity as the clinical signs often used to determine




Fig. 4. Evoked changes “total effect” in the striatal extracellular dialysate content of aspartate, glutamate and GABA following administration of kainic acid (12 mg/kg; solid bars) and SCH 23390 (0.5 mg/kg) + kainic acid (open bars), in the guinea pig. Results are expressed as a percent of the respective baseline values (mean \pm S.E.M., $n = 5$). ** $P < 0.01$; *** $P < 0.001$, kainic acid vs. SCH 23390 + kainic acid; two-tailed Mann–Whitney U -test.

stages of seizure activity, e.g., intermittent and status epilepticus, have been shown in certain cases not to relate to electrographical activity (13,18).

Studies in Other Diseases of the Central Nervous System

Parkinson's and Huntington's diseases

Blockade of the dopamine D_1 receptor subtype with a single dose of SCH 23390 increases the activity of aromatic L-amino acid decarboxylase (AAAD), which converts L-3,4-dihydroxyphenylalanine (L-DOPA) to dopamine. When L-DOPA is administered to rats after pretreatment with SCH 23390 there is a significant increase in the formation of DOPAC and dopamine turnover in striatum and midbrain compared with L-DOPA alone, suggesting further enhancement of dopamine metabolism. These studies suggest that it may be possible to enhance the conversion of L-DOPA to dopamine in Parkinson's disease patients by administering substances that augment brain AAAD, such as SCH 23390 (52). In Huntington's disease, studies with [^{11}C]SCH 23390 and [^{11}C]raclopride have shown that positron emission topography (PET) measures of human striatal D_1 and D_2 dopamine binding, respectively, can be used to identify asymptomatic Huntington's disease mutation carriers who are actively progressing and who would, thus, be suitable for putative neuroprotective therapies. Measures of disease progression rates in Huntington's disease patients and asymptomatic mutation carriers using these radiolabeled ligands will be of critical importance in future trials of experimental restorative treatments (7,33,72).




Fig. 5. Evoked changes “total effect” in the EEG power of both striatal and cortical EEG following administration of soman (30.5 µg/kg; solid bars) and SCH 23390 (0.5 mg/kg) + soman (open bars), in the guinea pig. Results are expressed as a percent of the respective baseline values (mean ± SEM, $n = 5$). ** $P < 0.01$; *** $P < 0.001$, soman vs. SCH 23390 + soman; two-tailed Mann–Whitney U -test.

Psychosis

The role of D_1 receptors in schizophrenia is tentative (23,69) but in preclinical trials SCH 23390 has been reported to have an antipsychotic action in schizophrenic patients, without inducing extrapyramidal side effects, such as spasms, restlessness, and pseudoparkinsonism (24,25). Hietala et al. (39) have, however, concluded from the majority of the available data that SCH 23390 does not share the profile of atypical neuroleptics. Farde (30) showed that administration of SCH 23390 to four healthy subjects at doses of 310–810 µg, i.v., induced akathisia, the syndrome of motor restlessness, at the three highest doses. However, akathisia is a transient effect (32). As with the prototypic atypical antipsychotic clozapine, that is routinely used in the treatment of schizophrenia, SCH 23390 may be exerting its direct/indirect action with nondopaminergic systems, such as 5-HT. Unfortunately, based on the current literature on the relevance of the D_1 receptor to higher cognition, it would be contradictory to block D_1 receptors in schizophrenia to relieve their cognitive deficits and associated psychotic symptoms. Indeed, recent clinical trials of the more selective D_1 receptor antagonist SCH 39166 revealed that it is incapable of improving schizophrenic symptoms (44). Moreover, recent electrophysiological findings of a D_1 potentiation of NMDA receptor functions in the PFC, striatum, and hippocampus may further illustrate the role of the D_1 receptor in improving cognition, as well as in inducing seizures as a result of the overstimulation of NMDA receptor (19,20,58).

Other idiopathic psychotic disorders in which dopamine receptors have been implicated include paranoia, mania, Tourette’s syndrome, major depression, childhood attention–hyperactivity disorder and substance abuse/addiction.




Fig. 6. Evoked changes “total effect” in the EEG power of both striatal and cortical EEG following administration of kainic acid (12 mg/kg; solid bars) and SCH 23390 (0.5 mg/kg) + kainic acid (open bars), in the guinea pig. Results are expressed as a percent of the respective baseline values (mean \pm S.E.M., $n = 5$). ** $P < 0.01$; *** $P < 0.001$, kainic acid vs. SCH 23390 + kainic acid; two-tailed Mann–Whitney U -test.

CONCLUSIONS

The anticonvulsant properties of SCH 23390 highlight the importance of the D_1 dopaminergic mechanisms in facilitating the initiation and spread of generalized seizures and that inhibition of the release-enhancing dopamine D_1 receptor located on the terminals of the GABAergic striatonigral neurons possibly attenuates seizure activity. This evidence is further supported by results from the rat showing that the D_1 agonist SKF 38393 (30 mg/kg, i.p.) is proconvulsant in a pilocarpine model of epileptiform activity by virtue of its activity at dopamine D_1 receptors (4,2,10). It has also been concluded that the normal synergistic balance of the dopamine D_1 and D_2 receptor subtype activity confers resistance to seizure-promoting stimuli (2). Further investigation with the selective antagonist SCH 23390, utilizing techniques to monitor neurochemistry and electrophysiology should yield a better mechanistic picture of the role of these receptors in many paradigms.

While there is no immediate prospect of developing D_1 receptor antagonists, specifically SCH 23390, as clinically useful antiepileptics, there is a growing awareness that seizures might be precipitated as a consequence of the treatment of other neurological disorders, e.g., the use of D_1 agonists in Parkinson's disease. Unfortunately, the pharmacokinetic properties of the drug and the results of previous animal studies (70), which indicate that the D_1 receptor antagonists can induce stereotypical and atypical behaviors, make SCH 23390 somewhat unlikely as a possible therapy. However, a better understanding of the involvement of striatal dopaminergic receptor subtypes, with respect to seizure ac-

tivity, has implications for these receptors as a possible target for the prevention or amelioration of epileptic seizures.

REFERENCES

1. Abekawa T, Ohmori T, Ito K, Koyama T. D₁ dopamine receptor activation reduces extracellular glutamate and GABA concentrations in the medial prefrontal cortex. *Brain Res* 2000;867:250–254.
2. al-Tajir G, Starr MS. Anticonvulsant action of SCH 23390 in the striatum of the rat. *Eur J Pharmacol* 1990; 191:329–336.
3. al-Tajir G, Starr MS. D₂ agonists protect rodents against pilocarpine-induced convulsions by stimulating D₂ receptors in the striatum, but not in the substantia nigra. *Pharmacol Biochem Behav* 1991;39:109–113.
4. al-Tajir G, Chandler CJ, Starr BS, Starr MS. Opposite effects of stimulation of D₁ and D₂ dopamine receptors on the expression of motor seizures in mouse and rat. *Neuropsychopharmacology* 1990;29:657–661.
5. Alam AM, Starr MS. Dopaminergic modulation of pilocarpine-induced motor seizures in the rat: The role of hippocampal dopamine D₁ receptors. *Eur J Pharmacol* 1992;222:227–232.
6. Anden NE, Dahlstrom A, Fuxe K, Larsson K. Mapping out of catecholamine and 5-hydroxytryptamine neurons innervating the telencephalon and diencephalon. *Life Sci* 1965;4:1275–1279.
7. Andrews TC, Weeks RA, Turjanski N, et al. Huntington's disease progression. PET and clinical observations. *Brain* 1999;122:2353–2363.
8. Arnsten AF, Cai JX, Murphy BL, Goldman-Rakic PS. Dopamine D₁ receptor mechanisms in the cognitive performance of young adult and aged monkeys. *Psychopharmacology (Berl)* 1994;116:143–151.
9. Barasi S, Ben-Sreti MM, Clatworthy AL, et al. Dopamine receptor-mediated spinal antinociception in the normal and haloperidol pretreated rat: Effects of sulpiride and SCH 23390. *Br J Pharmacol* 1987;90:15–22.
10. Barone P, Palma V, de Bartolomeis A, Cicarelli G, Campanella G. Dopaminergic regulation of epileptic activity. *Neurochem Int* 1992;20(Suppl):245S–249S.
11. Billard W, Ruperto V, Crosby G, Iorio LC, Barnett A. Characterization of the binding of ³H-SCH 23390, a selective D₁ receptor antagonist ligand, in rat striatum. *Life Sci* 1984;35:1885–1893.
12. Bischoff S, Heinrich M, Sonntag JM, Krauss J. The D₁ dopamine receptor antagonist SCH 23390 also interacts potently with brain serotonin (5-HT₂) receptors. *Eur J Pharmacol* 1986;129:367–370.
13. Bourne JA, Fosbraey P. Novel method of monitoring electroencephalography at the site of microdialysis during chemically evoked seizures in a freely moving animal. *J Neurosci Meth* 2000;99:85–90.
14. Bourne JA, Fosbraey P, Halliday J. SCH 23390 affords protection against soman-evoked seizures in the freely moving guinea-pig: A concomitant neurochemical, electrophysiological and behavioural study. *Neuropharmacology* 2001;40:279–288.
15. Bourne JA, Fosbraey P, Halliday J. Changes in striatal electroencephalography and neurochemistry induced by kainic acid seizures are modified by dopamine receptor antagonists. *Eur J Pharmacol* 2001;413: 189–198.
16. Burke K, Chandler CJ, Starr BS, Starr MS. Seizure promotion and protection by D₁ and D₂ dopaminergic drugs in the mouse. *Pharmacol Biochem Behav* 1990;36:729–733.
17. Carlsson A. On the neuronal circuitries and neurotransmitters involved in the control of locomotor activity. *J Neural Transm Suppl* 1993;40:1–12.
18. Carpentier P, Delamanche IS, Le Bert M, Blanchet G, Bouchaud C. Seizure-related opening of the blood-brain barrier induced by soman: Possible correlation with the acute neuropathology observed in poisoned rats. *Neurotoxicology* 1990;11:493–508.
19. Cepeda C, Colwell CS, Itri JN, Chandler SH, Levine MS. Dopaminergic modulation of NMDA-induced whole cell currents in neostriatal neurons in slices: Contribution of calcium conductances. *J Neurophysiol* 1998;79:82–94.
20. Cepeda C, Colwell CS, Itri JN, Gruen E, Levine MS. Dopaminergic modulation of early signs of excitotoxicity in visualized rat neostriatal neurons. *Eur J Neurosci* 1998;10:3491–3497.
21. Christensen AV, Arnt J, Hyttel J, Svendsen O. Behavioural correlates to the dopamine D₁ and D₂ antagonists. *Pol J Pharmacol Pharm* 1984;36:249–264.
22. Christensen AV, Arnt J, Hyttel J, Larsen JJ, Svendsen O. Pharmacological effects of a specific dopamine D₁ antagonist SCH 23390 in comparison with neuroleptics. *Life Sci* 1984;34:1529–1540.

23. Clark D, White FJ. D₁ dopamine receptor—the search for a function: A critical evaluation of the D₁/D₂ dopamine receptor classification and its functional implications. *Synapse* 1987;1:347–388.
24. Coffin VL, Latranyi MB, Chipkin RE. Acute extrapyramidal syndrome in Cebus monkeys: Development mediated by dopamine D₂ but not D₁ receptors. *J Pharmacol Exp Ther* 1989;249:769–774.
25. Creese I, Chen A. Selective D₁ dopamine receptor increase following chronic treatment with SCH 23390. *Eur J Pharmacol* 1985;109:127–128.
26. Cuomo V, Cagiano R, Colonna M, Renna G, Racagni G. Influence of SCH 23390, a DA1-receptor antagonist, on the behavioural responsiveness to small and large doses of apomorphine in rats. *Neuropharmacology* 1986;25:1297–1300.
27. Dalley JW, Stanford SC. Incremental changes in extracellular noradrenaline availability in the frontal cortex induced by naturalistic environmental stimuli: A microdialysis study in the freely moving rat. *J Neurochem* 1995;65:2644–2651.
28. Donnelly D, Findlay JB, Blundell TL. The evolution and structure of aminergic G protein-coupled receptors. *Receptors Channels* 1994;2:61–78.
29. Fage D, Scatton B. Opposing effects of D₁ and D₂ receptor antagonists on acetylcholine levels in the rat striatum. *Eur J Pharmacol* 1986;129:359–362.
30. Farde L. Selective D₁- and D₂-dopamine receptor blockade both induces akathisia in humans — a PET study with [¹¹C]SCH 23390 and [¹¹C]raclopride. *Psychopharmacology (Berl)* 1992;107:23–29.
31. Fosbraey P. Amphetamine-evoked dopamine release from guinea-pig striatum is potentiated by environmental stress: A need for habituation with microdialysis equipment. *Neuropharmacology* 1993;7:190–194.
32. Gessa GL, Canu A, Del Zompo M, Burrai C, Serra G. Lack of acute antipsychotic effect of SCH 23390, a selective dopamine D₁ receptor antagonist. *Lancet* 1991;337:854–855.
33. Ginovart N, Lundin A, Farde L, et al. PET study of the pre- and post-synaptic dopaminergic markers for the neurodegenerative process in Huntington's disease. *Brain* 1997;120:503–514.
34. Granon S, Passetti F, Thomas KL, Dalley JW, Everitt BJ, Robbins TW. Enhanced and impaired attentional performance after infusion of D₁ dopaminergic receptor agents into rat prefrontal cortex. *J Neurosci* 2000;20:1208–1215.
35. Hall H, Sedvall G, Magnusson O, Kopp J, Halldin C, Farde L. Distribution of D₁- and D₂-dopamine receptors, and dopamine and its metabolites in the human brain. *Neuropsychopharmacology* 1994;11:245–256.
36. Harsing LG, Zigmond MJ. Influence of dopamine on GABA release in striatum: Evidence for D₁-D₂ interactions and non-synaptic influences. *Neuroscience* 1997;77:419–429.
37. Hauber W. Blockade of subthalamic dopamine D₁ receptors elicits akinesia in rats. *Neuroreport* 1998;9:4115–4118.
38. Hauber W, Lutz S. Dopamine D₁ or D₂ receptor blockade in the globus pallidus produces akinesia in the rat. *Behav Brain Res* 1999;106:143–150.
39. Hietala J, Lappalainen J, Koulu M, Syvalahti E. Dopamine D₁ receptor antagonism in schizophrenia: Is there reduced risk of extrapyramidal side-effects? *Trends Pharmacol Sci* 1990;11:406–410.
40. Hyttel J. SCH 23390—the first selective dopamine D₁ antagonist. *Eur J Pharmacol* 1983; 91:153–154.
41. Iorio LC, Barnett A, Leitz FH, Houser VP, Korduba CA. SCH 23390, a potential benzazepine antipsychotic with unique interactions on dopaminergic systems. *J Pharmacol Exp Ther* 1983;226:462–468.
42. Itoh Y, Beaulieu M, Kebabian JW. The chemical basis for the blockade of the D-1 dopamine receptor by SCH 23390. *Eur J Pharmacol* 1984;100:119–122.
43. Jarvie KR, Caron MG. Heterogeneity of dopamine receptors. *Adv Neurol* 1993;60:325–333.
44. Karlsson P, Smith L, Farde L, Harnryd C, Sedvall G, Wiesel F-A. Lack of apparent antipsychotic effect of the D₁-dopamine receptor antagonist SCH 39166 in acutely ill schizophrenic patients. *Psychopharmacology (Berl)* 1995;121:309–316.
45. Kebabian JW, Calne DB. Multiple receptors for dopamine. *Nature* 1979;277:93–96.
46. Kilts CD, Dew KL, Ely TD, Mailman RB. Quantification of R-(+)-7-chloro-8-hydroxy-1-phenyl-2,3,4,5-tetrahydro-1H-3-methyl-3-benzazepine in brain and blood by use of reversed-phase high-performance liquid chromatography with electrochemical detection. *J Chromatogr* 1985;342:452–457.
47. Krebs MO, Trovero F, Desban M, Gauchy C, Glowinski J, Kemel ML. Distinct presynaptic regulation of dopamine release through NMDA receptors in striosome- and matrix-enriched areas of the rat striatum. *J Neurosci* 1991;11:1256–1262.
48. Krebs MO, Desce JM, Kemel ML, et al. Glutamatergic control of dopamine release in the rat striatum: Evidence for presynaptic N-methyl-D-aspartate receptors on dopaminergic nerve terminals. *J Neurochem* 1991;56:81–85.

49. Lachowicz JE, Sibley DR. Molecular characteristics of mammalian dopamine receptors. *Pharmacol Toxicol* 1997;81:105–113.
50. Morari M, O'Connor WT, Ungerstedt U, Fuxe K. Dopamine D₁ and D₂ receptor antagonism differentially modulates stimulation of striatal neurotransmitter levels by *N*-methyl-D-aspartic acid. *Eur J Pharmacol* 1994;256:23–30.
51. Murphy BL, Arnsten AF, Goldman-Rakic PS, Roth RH. Increased dopamine turnover in the prefrontal cortex impairs spatial working memory performance in rats and monkeys. *Proc Natl Acad Sci USA* 1996;93:1325–1329.
52. Neff NH, Wemlinger TA, Hadjiconstantinou M. SCH 23390 enhances exogenous L-DOPA decarboxylation in nigrostriatal neurons. *J Neural Transm* 2000;107:429–443.
53. O'Boyle KM, Waddington JL. Identification of the enantiomers of SK&F 83566 as specific and stereoselective antagonists at the striatal D₁ dopamine receptor: Comparisons with the D-2 enantioselectivity of Ro 22-1319. *Eur J Pharmacol* 1984;106:219–220.
54. Pettersson I, Gundertofte K, Palm J, Liljefors T. A study on the contribution of the 1-phenyl substituent to the molecular electrostatic potentials of some benzazepines in relation to selective dopamine D₁ receptor activity. *J Med Chem* 1992;35:502–507.
55. Previc FH. Dopamine and the origins of human intelligence. *Brain Cogn* 1999;41:299–350.
56. Ribak C. GABA and the basal ganglia. In: *The GABAergic Neurons of the Extrapyramidal System as Revealed by Immunohistochemistry*. DiChiara C, Gessa G, Eds. New York: Raven Press, 1981:23–36.
57. Santiago M, Westerink BH. Characterization and pharmacological responsiveness of dopamine release recorded by microdialysis in the substantia nigra of conscious rats. *J Neurochem* 1991;57:738–747.
58. Seamans JK, Durstewitz D, Christie BR, Stevens CF, Sejnowski TJ. Dopamine D₁/D₅ receptor modulation of excitatory synaptic inputs to layer V prefrontal cortex neurons. *Proc Natl Acad Sci USA* 2001;98:301–306.
59. Sokoloff P, Schwartz JC. Novel dopamine receptors half a decade later. *Trends Pharmacol Sci* 1995;16:270–275.
60. Sokoloff P, Giros B, Martres MP, Bouthenet ML, Schwartz JC. Molecular cloning and characterization of a novel dopamine receptor (D₃) as a target for neuroleptics. *Nature* 1990;347:146–151.
61. Starr BS, Starr MS. Behavioural interactions involving D₁ and D₂ dopamine receptors in non-habituated mice. *Neuropharmacology* 1987;26:613–619.
62. Starr M. Opposing roles of dopamine D₁ and D₂ receptors in nigral γ -[³H]aminobutyric acid release? *J Neurochem* 1987;49:1042–1049.
63. Starr MS. The role of dopamine in epilepsy. *Synapse* 1996;22:159–194.
64. Sunahara RK, Guan HC, O'Dowd BF, et al. Cloning of the gene for a human dopamine D₅ receptor with higher affinity for dopamine than D₁. *Nature* 1991;350:614–619.
65. Tiberi M, Caron MG. High agonist-independent activity is a distinguishing feature of the dopamine D_{1B} receptor subtype. *J Biol Chem* 1994;269:27925–27931.
66. Tomiyama K, Koshikawa N, Funada K, Oka K, Kobayashi M. *In vivo* microdialysis evidence for transient dopamine release by benzazepines in rat striatum. *J Neurochem* 1995;65:2790–2795.
67. Turski L, Cavalheiro EA, Czuczwar SJ, Turski WA, Kleinrok Z. The seizures induced by pilocarpine: Behavioral, electroencephalographic and neuropathological studies in rodents. *Pol J Pharmacol Pharm* 1987;39:545–555.
68. Van Tol HH, Bunzow JR, Guan HC, et al. Cloning of the gene for a human dopamine D₄ receptor with high affinity for the antipsychotic clozapine. *Nature* 1991;350:610–614.
69. Waddington JL. Therapeutic potential of selective D₁ dopamine receptor agonists and antagonists in psychiatry and neurology. *Gen Pharmacol* 1988;19:55–60.
70. Waddington JL, O'Boyle KM. Drugs acting on brain dopamine receptors: A conceptual re-evaluation five years after the first selective D₁ antagonist. *Pharmacol Ther* 1989;43:1–52.
71. Waddington JL, Molloy AG, O'Boyle KM, Mashurano M. Motor consequences of D-1 dopamine receptor stimulation and blockade. *Clin Neuropharmacol* 1986;9(Suppl 4):20–22.
72. Weeks RA, Piccini P, Harding AE, Brooks DJ. Striatal D₁ and D₂ dopamine receptor loss in asymptomatic mutation carriers of Huntington's disease. *Ann Neurol* 1996;40:49–54.
73. Westerink BH. Brain microdialysis and its application for the study of animal behaviour. *Behav Brain Res* 1995;70:103–124.

74. Westerink BH, de Vries JB. On the mechanism of neuroleptic induced increase in striatal dopamine release: Brain dialysis provides direct evidence for mediation by autoreceptors localized on nerve terminals. *Neurosci Lett* 1989;99:197–202.
75. Williams GV, Goldman-Rakic PS. Modulation of memory fields by dopamine D₁ receptors in prefrontal cortex. *Nature* 1995;376:572–575.
76. Yang CR, Seamans JK. Dopamine D₁ receptor actions in layers V-VI rat prefrontal cortex neurons *in vitro*: Modulation of dendritic-somatic signal integration. *J Neurosci* 1996;16:1922–1935.
77. Zetterstrom T, Sharp T, Ungerstedt U. Effect of dopamine D₁ and D₂ receptor selective drugs on dopamine release and metabolism in rat striatum *in vivo*. *Naunyn Schmiedeberg's Arch Pharmacol* 1986;334:117–124.